

THE VOLUME FLUX GROUP AND NONPOSITIVE CURVATURE

by Pablo Suárez-Serrato

ABSTRACT. We show that every closed nonpositively curved manifold with non-trivial volume flux group has zero minimal volume, and admits a finite covering with circle actions whose orbits are homologically essential. This proves a conjecture of Kedra–Kotschick–Morita for this class of manifolds.

Let M be a closed smooth manifold and μ a volume form on M . Denote by Diff^μ the group of μ -preserving diffeomorphisms of M , and by Diff_0^μ its identity component. The μ -flux homomorphism Flux_μ , from the universal covering $\widetilde{\text{Diff}}_0^\mu$ to the $(n-1)$ -cohomology group $H^{n-1}(M; \mathbf{R})$, is defined by the formula

$$\text{Flux}_\mu([\varphi_t]) = \int_0^1 [i_{\dot{\varphi}_t} \mu] dt.$$

It induces a homomorphism

$$\text{Flux}_\mu: \pi_1(\text{Diff}_0^\mu) \rightarrow H^{n-1}(M; \mathbf{R})$$

whose image is the volume flux group $\Gamma_\mu \subset H^{n-1}(M; \mathbf{R})$. The μ -flux homomorphism descends to a homomorphism

$$\text{Flux}_\mu: \text{Diff}_0^\mu \rightarrow H^{n-1}(M; \mathbf{R})/\Gamma_\mu.$$

For a closed connected smooth Riemannian manifold (M, g) , let $\text{Vol}(M, g)$ denote the volume of g and let K_g be its sectional curvature. We define the minimal volume of M following J Cheeger and M Gromov [2]:

$$\text{MinVol}(M) := \inf_g \{ \text{Vol}(M, g) : |K_g| \leq 1 \}$$

Date: November 12, 2008.

2000 Mathematics Subject Classification. Primary 57R50, 57S05, 53C23; Secondary 53C21, 22E65.

Key words and phrases. Volume flux group, minimal volume, nonpositive curvature.

The author is supported by the Deutsche Forschungsgemeinschaft (DFG) project ‘Asymptotic Invariants of Manifolds’.

The minimal volume is a very sensitive invariant, it was first observed by L Bessi eres [1] that its value may depend on the differentiable structure of (M, g) . Indeed, D Kotschick [6] has shown that even the vanishing of $\text{MinVol}(M)$ can detect changes in the smooth structure of M .

The investigation of relationships between the volume flux group of M and various invariants which bound $\text{MinVol}(M)$ from below was set in motion by J Kedra, D Kotschick and S Morita [5]. They put forward the idea that if a closed manifold has non-trivial volume flux group then its minimal volume should vanish. The aim of this note is to verify that statement for closed manifolds which carry a metric of nonpositive sectional curvature.

The volume flux group Γ_μ is independent of the form μ —see section 3 of [5]—so it can be considered as an invariant of the manifold M itself.

Theorem 1. *Every closed nonpositively curved manifold with non-trivial volume flux group has zero minimal volume.*

The technique we will use to show that the minimal volume vanishes is an $\mathcal{F}$ –structure, which was introduced by Gromov as a generalisation of an S^1 –action.

Definition 2. *An $\mathcal{F}$ –structure on a closed manifold M is given by the following conditions.*

- (1) *A finite open cover $\{U_1, \dots, U_N\}$*
- (2) *$\pi_i: \tilde{U}_i \rightarrow U_i$ a finite Galois covering with group of deck transformations Γ_i , $1 \leq i \leq N$*
- (3) *A smooth torus action with finite kernel of the k_i –dimensional torus,*
 $\phi_i: T^{k_i} \rightarrow \text{Diff}(\tilde{U}_i)$, $1 \leq i \leq N$
- (4) *A homomorphism $\Psi_i: \Gamma_i \rightarrow \text{Aut}(T^{k_i})$ such that*

$$\gamma(\phi_i(t)(x)) = \phi_i(\Psi_i(\gamma)(t))(\gamma x)$$

for all $\gamma \in \Gamma_i$, $t \in T^{k_i}$ and $x \in \tilde{U}_i$

- (5) *For any finite sub-collection $\{U_{i_1}, \dots, U_{i_l}\}$ such that $U_{i_1 \dots i_l} := U_{i_1} \cap \dots \cap U_{i_l} \neq \emptyset$ the following compatibility condition holds: let $\tilde{U}_{i_1 \dots i_l}$ be the set of points $(x_{i_1}, \dots, x_{i_l}) \in \tilde{U}_{i_1} \times \dots \times \tilde{U}_{i_l}$ such that $\pi_{i_1}(x_{i_1}) = \dots = \pi_{i_l}(x_{i_l})$. The set $\tilde{U}_{i_1 \dots i_l}$ covers $\pi_{i_j}^{-1}(U_{i_1 \dots i_l}) \subset \tilde{U}_{i_j}$ for all $1 \leq j \leq l$, then we require that ϕ_{i_j} leaves $\pi_{i_j}^{-1}(U_{i_1 \dots i_l})$ invariant and it lifts to an action on $\tilde{U}_{i_1 \dots i_l}$ such that all lifted actions commute*

An $\mathcal{F}$ -structure is said to be *pure* if all the orbits of all actions at a point, for every point have the same dimension. We will say an $\mathcal{F}$ -structure is *polarised* if the smooth torus action ϕ_i above are fixed point free for every U_i . The existence of a polarised $\mathcal{F}$ -structure on a manifold M implies the minimal volume $\text{MinVol}(M)$ is zero by the main result of Cheeger and Gromov [2], the interested reader is invited to consult the illuminating examples found there as well.

The attentive reader will notice that the definition of an $\mathcal{F}$ -structure above is different from the sophisticated one found in [2]. Despite this, it is sufficiently practical to be implemented and also satisfies the properties needed in the proof that a polarised $\mathcal{F}$ -structure forces the minimal volume to vanish, which can be consulted in [2].

Proof of Theorem 1. Let M be a compact nonpositively curved manifold whose volume flux group is not trivial. The fundamental group $\pi_1(M)$ has non-trivial centre Z , a proof can be found in [5, Theorem 15] and compared with work of A Fathi [4, Proposition 5.1] and the various references and attributions contained therein. In this case M admits a finite covering space M^* diffeomorphic to $T^k \times N$, where T^k is a flat torus of dimension k and N is a compact nonpositively curved manifold as was shown by P Eberlein [3].

Even though it may seem that this already implies $\text{MinVol}(M) = 0$, for completeness will now show how to construct a pure polarised $\mathcal{F}$ -structure on M ; since the torus T^k splits off M^* smoothly and the action of T^k on itself as a factor of M^* is compatible with the covering transformation in the required sense. This will also provide an example of a detailed construction of a pure polarised $\mathcal{F}$ -structure.

Represent M as H/Γ , where H is simply connected and Γ is a properly discontinuous group of isometries of H which acts freely. The space H decomposes into $H_1 \times H_2$, where $H_1 = \mathbf{R}^k$ is a Euclidean space of dimension $k = \text{rank}(Z)$.

Notice that every element γ of Γ is of the form $\gamma = \gamma_1 \times \gamma_2 \in \text{Iso}(H_1) \times \text{Iso}(H_2)$, here $\text{Iso}(H_i)$ denotes the group of isometries of H_i (see Lemma 1 in [3], and the subsequent discussion). Let $p_i: \Gamma \rightarrow \text{Iso}(H_i)$ denote the projection homomorphisms, then $\Gamma_1 = p_1(\Gamma)$ acts by translations on H_1 and $\Gamma_2 = p_2(\Gamma)$ is a discrete subgroup of $\text{Iso}(H_2)$. So $Z \subset \Gamma_1$ and H_1/Z is a compact flat torus T^k . The projection $p: H_1 \rightarrow T^k$ allows us to define $\rho: \Gamma_2 \rightarrow T^k$ by setting $\rho(p_2\gamma) = p(p_1\gamma)$. The function ρ is well defined since $\ker(p) = \ker(p_2) = Z$. The centre Z can also be thought of as a set of vectors in H_1 , as $Z \subset \Gamma_1$. In this way Z acts on H_1 by translations.

Recall that Γ_2 has trivial centre and that M is isometric to $(T^k \times H_2)/\Gamma_2$ [3], here Γ_2 acts on $T^k \times H_2$ by $\psi(s, h) = (\rho(\psi)s, \psi(h))$ with (s, h) in $T^k \times H_2$ and ψ in Γ_2 . This can be read from the diagram

$$\begin{array}{ccc} H = H_1 \times H_2 & \xrightarrow{p \times Id} & T^k \times H_2 \\ \downarrow & & \downarrow q \\ M = H/\Gamma & \xrightarrow{F} & (T^k \times H_2)/\Gamma_2 \end{array}$$

where F is defined so that the diagram commutes.

In Γ_2 there exists a finite index subgroup Γ_0 —denoted by Γ_2^{**} in [3]—which makes the following diagram commute.

$$\begin{array}{ccc} H & \xrightarrow{\quad} & T^k \times (H_2/\Gamma_0) := M^* \\ & \searrow & \downarrow \\ & & (T^k \times H_2)/\Gamma_2 = M \end{array}$$

Define $q^*: T^k \times (H_2/\Gamma_0) = M^* \rightarrow M$ as in the previous diagram and denote by Γ^* the group of deck transformations of q^* seen as a covering map. Notice that $\Gamma^* \subset \Gamma_2$.

The function ρ is defined on all of Γ_2 , so we can also consider the restriction of ρ to Γ^* .

We are now in a position to verify that this construction gives M a pure polarised $\mathcal{F}$ -structure, and hence $\text{MinVol}(M) = 0$ as claimed. Let us check that every condition which guarantees the existence of a polarised $\mathcal{F}$ -structure is met.

- (1) Take $U = M$, as an open cover with a single set.
- (2) Define $\tilde{U} := M^* = T^k \times N$, here $N = H_2/\Gamma_0$. The quotient map

$$q^*: \tilde{U} = T^k \times N \rightarrow (T^k \times H_2)/\Gamma_2 = M = U$$

is a finite Galois covering with deck transformation group Γ^* .

- (3) The k -torus T^k acts smoothly and without fixed points on itself as a factor of $T^k \times (H_2/\Gamma_0)$. So we have the action

$$\phi: T^k \rightarrow \text{Diff}(\tilde{U}) = \text{Diff}(T^k \times N)$$

given by $\phi(t)(s, h) = (s + t, h)$.

- (4) We will use the function ρ to define the automorphism $\Psi: \Gamma^* \rightarrow \text{Aut}(T^k)$, set $\Psi(\gamma) := \rho(\gamma)$. Let $t \in T^k$ and $x \in \tilde{U} = T^k \times N \Rightarrow x = (s, h) \in T^k \times N$. Notice that for s and t in T^k we have that $\rho(\gamma)(s + t) = (\rho(\gamma)s + \rho(\gamma)t)$ since $\rho(\gamma) \in T^k$.

We plug in this information to obtain the following equalities.

$$\begin{aligned}
 \gamma(\phi(t)x) &= \gamma(\phi(t)(s, h)) \\
 &= \gamma(s + t, h) \\
 &= (\rho(\gamma)(s + t), \gamma h) \\
 &= (\rho(\gamma)s + \rho(\gamma)t, \gamma h) = \star
 \end{aligned}$$

$$\begin{aligned}
 \phi(\Psi(\gamma)t)(\gamma x) &= \phi(\rho(\gamma)t)(\gamma x) \\
 &= \phi(\rho(\gamma)t)(\rho(\gamma)s, \gamma h) \\
 &= (\rho(\gamma)s + \rho(\gamma)t, \gamma h) = \star
 \end{aligned}$$

Therefore $\gamma(\phi_i(t)(x)) = \phi_i(\Psi_i(\gamma)(t))(\gamma x)$ and the condition is satisfied.

- (5) This condition does not need to be verified, because we only have one covering set.

Since the action of T^k on itself as a factor of $T^k \times (H_2/\Gamma_0)$ is fixed point free, the above construction gives M a pure polarised $\mathcal{F}$ -structure. $\square$

We will state the contrapositive statement to the theorem because it strengthens Corollary 17 of [5]: *Let M be a closed nonpositively curved manifold with positive minimal volume. Then the volume flux group Γ_μ is trivial for every volume form μ on M .*

Another noteworthy feature of the covering $M^* \rightarrow M$ is that on $M^* \cong T^k \times N$ elements of the volume flux group which come from rotations on the T^k factor are circle actions with homologically essential orbits. It is not yet clear if in the general case a non-trivial volume flux group implies existence of a circle action with homologically essential orbits at least in a multiple cover—compare with Remark 19 in [5]—but it is virtually so for every closed nonpositively curved manifold M , since it is true for M^* .

Acknowledgements: I wish to warmly thank Dieter Kotschick for a number of interesting conversations and for commenting on a previous version of this work.

REFERENCES

- [1] L. Bessières, *Un théorème de rigidité différentielle*, Comment. Math. Helv. 73 (1998), no. 3, 443–479.
- [2] J. Cheeger and M. Gromov, *Collapsing Riemannian Manifolds while keeping their curvature bounded I*, Jour.Differential Geom. 23 (1986) 309–346.

- [3] P. Eberlein, *A canonical form for compact nonpositively curved manifolds whose fundamental groups have nontrivial center*, Math. Ann. 260 (1982) no. 1, 23–29.
- [4] A. Fathi, *Structure of the group of homeomorphisms preserving a good measure on a compact manifold*, Ann. Sci. École Norm. Sup. (4) 13 (1980), no. 1, 45–93.
- [5] J. Kedra, D. Kotschick and S. Morita, *Crossed Flux Homomorphisms and Vanishing Theorems for Flux Groups*, GAFA 16 (2006) 1246–1273.
- [6] D. Kotschick, *Entropies, Volumes and Einstein Metrics*, Preprint (2004) [arXiv:math.DG/0410215](https://arxiv.org/abs/math/0410215).

MATHEMATISCHES INSTITUT LMU, THERESIENSTRASSE 39, MÜNCHEN 80333 GERMANY.

CIMAT, CP: 36240, GUANAJUATO, GTO, MÉXICO.
E-mail address: p.suarez-serrato@cantab.net